lockstep with oxygen, as expected for primary elements. The fact that the nitrogen abundance gradient is not much steeper suggests that much of the nitrogen may also be of primary origin. A further puzzle arises in the fact that several isotopic ratios (¹⁸O/¹⁷O, C/¹³C, S/³⁴S, N/¹⁵N, etc.), measured at millimeter wavelengths, are constant to a high degree over the plane of




Fig. 3: Variations of electron temperature and preliminary abundance ratios He⁺/H⁺, O/H, and N/H, as a function of distance from the galactic centre. Arrows to the right indicate values for the LMC (30 Doradus) and SMC (N66).

the galaxy, in apparent contradiction to the marked gradients in O/H, S/H, etc. (Wannier, 1980, *Ann. Rev. Astron. Astrophys.* **18**, 399). These facts seem to call for a revision of our ideas about nucleosynthesis.

It is thought that the disk of our galaxy formed gradually, with infall of primordial gas extending over a long period. The main evidence for this is the shortage of old stars in the disk with low abundances. These infall models share the prediction that the abundance gradient should flatten off in the inner regions of the disk (Tinsley and Larson, 1978, *Astrophys. J.* 221, 554; Chiosi, 1980, *Astron. Astrophys.* 83, 206). There is no evidence for this in Fig. 3, but there are a number of ways out of this dilemma, such as postulating infall of metal-enriched gas from stars in the galactic bulge. There are clearly many free parameters in such models, but an increasing array of observational data will hopefully provide the constraints necessary to ultimately distinguish the actual evolutionary scenario.

List of Preprints Published at ESO Scientific Group

December 1981 - February 1982

- J. Breysacher, A.F.J. Moffat and V.S. Niemela: Wolf-Rayet Stars in the Magellanic Clouds. II. The Peculiar Eclipsing Binary HD 5980 in the SMC. Astrophysical Journal. December 1981.
- 178. E. Athanassoula, A. Bosma, M. Crézé and M.P. Schwarz: On the Sizes of Rings and Lenses in Disk Galaxies. Astronomy and Astrophysics. December 1981.
- I.S. Glass, A.F.M. Moorwood and W. Eichendorf: Mid-Infrared Observations of Seyfert 1 and Narrow-Line X-ray Galaxies. Astronomy and Astrophysics. December 1981.
- S. D'Odorico and M. Rosa; Wolf-Rayet Stars Associated to Giant Regions of Star Formation. IAU Symposium 99 on Wolf-Rayet Stars. December 1981.
- J. Materne, G. Chincarini, M. Tarenghi and U. Hopp: Optical Investigations of Two X-ray Clusters of Galaxies: 0430.6-6133 and 0626.7-5426. Astronomy and Astrophysics. January 1982.
- M. Rosa and S. D'Odorico: Wolf-Rayet Stars in Extragalactic H II Regions: II. NGC 604—A Giant H II Region Dominated by many Wolf-Rayet Stars. Astronomy and Astrophysics. January 1982.
- W. Bijleveld and E.A. Valentijn: Radio and X-ray Observations of the Abell 2241 Galaxy Clusters. Astronomy and Astrophysics. January 1982.
- Ch. Motch, J. van Paradijs, H. Pedersen, S.A. Ilovaisky und C. Chevalier: Visual and Infrared Photometry of 2A 0311-227. Astronomy and Astrophysics. January 1982.
- 185. L. Woltjer and G. Setti: Quasars in the Universe. To be published in the Proceedings of the Vatican Study Week on "Cosmology and Fundamental Physics", Rome, Sept. 28–Oct. 2, 1981. February 1982.
- 186. G. Setti and L. Woltjer: The Origin of the X- and γ-ray Backgrounds. To be published in the Proceedings of the Vatican Study Week on "Cosmology and Fundamental Physics", Rome, Sept. 28–Oct. 2, 1981. February 1982.
- R. Barbon, M. Capaccioli, R.M. West and R. Barbier: Redshifts of Parent Galaxies of Supernovae. Astronomy and Astrophysics, Suppl. February 1982.
- 188. R.M. West and H.E. Schuster: The ESO Quick Blue Survey and ESO (B) Atlas. Astronomy and Astrophysics, Suppl. February 1982.
- W. Eichendorf, A. Heck, B. Caccin, G. Russo and C. Sollazzo: UV, Optical and IR Observations of the Cepheid R. Muscae. Astronomy and Astrophysics. February 1982.
- 190. M.P. Véron-Cetty, P. Véron and M. Tarenghi: Are All Galactic Nuclear Regions Sodium Rich? Astronomy and Astrophysics. Februar 1982.