Leader, and R. Mehlman, "Hot spots on lo: Initial results from Galileo's near infrared mapping spectrometer", *Geophys. Res. Lett.* **24**, 2439 (1997).

- [27] J. R. Spencer, J. A. Stansberry, C. Dumas, D. Vakil, R. Pregler, M. Hicks, and K. Hege, "History of high-temperature lo volcanism: February 1995 to May 1997", *Geophys. Res. Lett.* 24, 2451 (1997).
- [28] F. Marchis, R. Prangé, and J.C. Christou, "Adaptive Optics mapping of

lo's volcanism in the thermal IR", *Icarus* (1999), submitted.

- [29] D. Le Mignant, E. Gendron, and F. Marchis, "The ESO ADONIS adaptive optics system: Study of the 3-5μm background emissivity", in ESO conference and workshop proceedings 56, 575 (Garching, Germany, 1998).
- [30] H. Geoffray, "Thermal background blurring on ADONIS system", in ESO conference and workshop proceedings 56, 531 (Garching, Germany, 1998).
- [31] R. Prangé, F. Marchis, and T. Fusco, "Mapping and monitoring of lo's hot spots by use of the ESO Adaptive Optics system.", in *31st Annual Meeting of the DPS*, (Padova, Italy, 1999).
- [32] F. Marchis, J. Berthier, P. Descamps, T. Fusco, R. Prangé, and T. Sekiguchi, "Ground-based high resolution observations of the Uranian system in the near IR", in *31st Annual Meeting of the DPS*, (Padova, Italy, 1999).

A New Look at the Sombrero Galaxy



This image of Messier 104, also known as Sombrero Galaxy, because of its particular shape, was obtained with FORS1 at VLT Antu on January 30, 2000. The colour image was made by a combination of three CCD images obtained by Peter Barthel from the Kapteyn Institute at Groningen, Netherlands. He and Mark Neeser, also from the Kapteyn Institute, produced the composite images. The "Sombrero" is notable for its dominant nuclear bulge, composed primarily of old stars, and its nearly edge-on disk composed of stars,

The "Sombrero" is notable for its dominant nuclear bulge, composed primarily of old stars, and its nearly edge-on disk composed of stars, gas and intricately structured dust. The complexity of this dust and the high resolution of this image are most apparent directly in front of the bright nucleus, but it is also evident as dark absorbing lanes throughout the disk.