

The *Mid*-Infrared  
*Interferometric*  
Instrument for the  
Very Large Telescope  
Interferometer

**MIDI**

—

**MIDI Guaranteed Time Observations  
Abstracts and Source List for April 2008 - Period 82**

Date 31 January 2008

### Summary of the guaranteed time programme for MIDI

Coordinator	Topic	Obs. Time UTs	Time ATs	Page
Klaus Meisenheimer	Dust Tori in Nearby Active Galactic Nuclei	65 h	–	<b>done</b>
Christoph Leinert	Studying the inner disks of low-mass young stellar objects	65 h	90 h	1
Anne Dutrey	Studying Inner Disks around Intermediate-mass Pre-Main-Sequence and Vega-type stars	62.5 h	100 h	3
Markus Feldt	A detailed study of massive young stars	52.5 h	305 h	5
Olivier Chesneau	The dusty circumstellar environment of Hot Stars	2 h	68 h	7
Andreas Quirrenbach	Cool Late Type Stars and related objects	25 h	450 h	9
Bruno Lopez	Extra-solar planets and brown dwarfs	25 h	–	11
	Allowance for rounding errors	3 h		
<b>TOTAL</b>		<b>300 h</b>	<b>110 n</b>	

**MIDI guaranteed time observations –  
Abstract and source list**

**Studying the inner disks  
of low-mass young stellar objects**

Authors: Ch. Leinert (coordinator), A. Dutrey, Th. Henning, T. Herbst, S. Ligi, B. Lopez,  
I. Pascucci, F. Przygodda, G. Niccolini, Th. Ratzka, H. Zinnecker

## **1 Abstract**

Using the spatial resolution of MIDI on the VLTI, about 2 AU at the 150 pc distance of nearby star forming regions, we want to perform a **direct** study of the properties of disks around T Tauri stars. The immediate outcome of these observations should be 10  $\mu\text{m}$  size, geometry and orientation of these disks, supplemented for the brighter objects by some information on chemical composition of dust and on physical structure and peculiarities in spatial distribution of circumstellar material. The latter effect may point to dust destruction or to depletion as it might result from gravitational interaction with an orbiting companion or planet. The results are relevant for the formation, structure and evolution of circumstellar disks which connect the formation of low-mass stars to the possible formation of planets.

For this study, the proposal contains a number of classical T Tauri stars of different mass and age. Binary stars are included: they allow us to look for differential effects and to study the results of dynamical interaction. Object groups known for particular accretion activity, like FUORs and binaries containing so-called infrared companions, are well represented on purpose – the emphasis is on the earlier phases of circumstellar disk evolution.

Total observing time : **65 hours on UTs - used up**  
**90 hours on ATs**

## 2 Source list

⇒ these eight objects are protected for AT observing only

apparently single stars								
target name	coordinates ( $\alpha, \delta$ ) (J2000)						telescopes	priority <sup>a</sup>
RY Tau	04	21	57.4	+28	26	35	ATs	***
DG Tau	04	27	04.7	+26	06	17	ATs	***

<sup>a</sup>highest priority is indicated by \*\*\*

multiple stars								
target name	coordinates ( $\alpha, \delta$ ) (J2000)						telescopes	priority <sup>a</sup>
T Tau	04	21	59.4	+19	32	06	ATs	***
Haro 6-10	04	29	24.4	+24	33	02	ATs	***
Z Cma	05	45	22.6	+09	04	12	ATs	***
Glass I	11	08	16.0	-77	33	47	ATs	***
AS 205	16	11	31.4	-18	38	25	ATs	**
VV CrA	19	03	06.7	-37	12	51	ATs	***

<sup>a</sup>highest priority is indicated by \*\*\*

**MIDI guaranteed time observations –  
Abstract and source list**

**Studying Inner Disks around  
Intermediate-mass YSOs and Vega-type stars**

Authors: Anne Dutrey (coordinator, Grenoble), Roy van Boekel (Amsterdam), Thomas Henning (Heidelberg), Christoph Leinert (Heidelberg), Bruno Lopez (Nice), Gilles Niccolini (Nice), Bringfried Stecklum (Tautenburg), Rens Waters (Amsterdam)

## **1 Abstract**

We propose to use MIDI at the VLTI to observe the dust disks surrounding nearby isolated Herbig Ae/Be stars and Vega-type stars ( $D \leq 450$  pc). **The spatial resolution achieved by MIDI (about 2 AU at 150 pc) will allow us to study *in-situ* the size at  $10 \mu\text{m}$  and structure of the inner regions of the proto-planetary disks surrounding Herbig Ae/Be stars and to search for warm dust in Vega-type systems.** These observations will put strong constraints on the models of disks surrounding intermediate mass young stars ( $2 - 5 M_{\odot}$ ). The immediate outcome should be the  $10 \mu\text{m}$  size, geometry, and orientation of disks. For a few bright objects, we also expect information on the spatial distribution of the carbonaceous and silicate dust components, that show strong spectral features in the  $10 \mu\text{m}$  region. The sample contains a number of stars of different mass and age. We have included single stars and binaries to study the effect of multiplicity on disk structure; the sample also contains some main sequence stars with dust shells, which allow us to investigate the evolution from gas-rich accretion disks towards gas-poor collisionally evolving systems.

This proposal should be considered as the intermediate-mass counterpart of the proposal on TTauri disks.

Total observing time : **62.5 hours on UTs – used up**  
**100 hours on ATs**

## 2 Source list

⇒ these ten objects are protected for AT observing only

apparently single stars								
target name	coordinates ( $\alpha, \delta$ ) (J2000)						telescopes	priority
MWC 758	05	30	27.53	+25	19	57.1	ATs	***
HD 41511	06	04	59.09	-16	29	03.6	ATs	**
HD 97048	11	08	04.6	-77	39	17	ATs	**
HD 100453	11	33	05.73	-54	19	27.8	ATs	***
HD 100546	11	33	25.64	-70	11	41.6	ATs	***
HD 104237	12	00	05.54	-78	11	32.5	ATs	***
HD 142527	15	56	41.86	-42	19	21.5	ATs	***
HD 163296	17	56	21.26	-21	57	19.5	ATs	***

multiple stars								
target name	coordinates ( $\alpha, \delta$ ) (J2000)						telescopes	priority
HD 150193	16	40	17.83	-23	53	44.1	ATs	*** (1.'07 - 2p)
TY CRA	19	01	40.83	-36	52	33.9	ATs	** (eclip/sb2 - 1p)

“2p” means separate pointings for the two components, “1p” means one combined pointing

**MIDI guaranteed time observations –  
Abstract and source list**

**A detailed study of massive young stars**

M. Feldt<sup>1</sup> (coordinator), Th. Henning<sup>1</sup>, L. Kaper<sup>2</sup>, Ch. Leinert<sup>1</sup>, H. Linz<sup>3</sup>, I. Pascucci<sup>1</sup>,  
M. Robberto<sup>4</sup>, B. Stecklum<sup>3</sup>, R. Waters<sup>2</sup>, H. Zinnecker<sup>5</sup>

<sup>1</sup> Max–Planck–Institut für Astronomie (MPIA) Heidelberg

<sup>2</sup> Astronomical Institute, University of Amsterdam

<sup>3</sup> Thüringer Landessternwarte Tautenburg

<sup>4</sup> ESA–Space Telescope Science Institute (STScI) Baltimore

<sup>5</sup> Astrophysikalisches Institut Postdam

**1 Abstract**

We propose to use MIDI to spatially resolve the circumstellar environment of prominent high-mass protostars and young stars. The dust distribution in the immediate stellar surroundings will allow us to constrain the evolutionary status of the sources. According to the standard scenario, the mid-IR emission is expected to be roughly symmetric in the early radial infall phases, asymmetric in the main disk-accretion phases, absent in the final stages with an established wind-cleared cavity and compact HII region. An enhanced fraction of close companions would provide support to the alternative hypothesis that massive stars form by coalescence of lower mass objects in the densest parts of young clusters.

Total observing time : **52.5 hours on UTs**  
**305 hours on ATs**

## 2 Source list

⇒ nine of the following objects are planned for UT observing, twelve are planned for observations on ATs, and four are planned for observations with both, UTs and ATs (see column “UT/AT” in the table)

Target	R.A. (J2000)	Dec. (J2000)	Age group / question class <sup>a</sup>	UT/AT
G045.45+00.06 MIR	19 14 21.17	+11 09 15.6	0	UT
G010.8399–02.5922	18 19 12.12	–20 47 29.7	I	UT
G284.01–0.85 #1	10 20 15.09	–58 03 39	I	UT + AT <sup>b</sup>
G268.4217–00.8486	09 01 54.4	–47 44 11	I	UT + AT <sup>b</sup>
Mon R2–IRS 2	06 07 45.8	–06 22 53.7	I	AT
Mon R2–IRS 3 NE	06 07 47.8	–06 22 55.7	I	UT
Mon R2–IRS 3 SW	06 07 47.8	–06 22 56.6	I	AT
M 8E IR	18 04 53.3	–24 26 42	I	AT
AFGL 4176	13 43 02.1	–62 08 52	I	AT
G345.0080+01.7946	16 56 47.6	–40 14 25	0/I	UT
G009.6164+00.1965	18 06 14.78	–20 31 40.5	I	UT
G305.2017+00.2071	13 11 10.5	–62 34 39	I	AT
Orion IRc2	05 35 14.4	–05 22 29.5	I	UT
AFGL 2136	18 22 26.5	–13 30 12	I	UT + AT <sup>b</sup>
G203.3167+02.0560	06 41 10.10	+09 29 34.0	I	AT
G324.2+0.12	15 32 54.0	–55 55 00	I	UT
G318.9491–00.1967	15 00 55.4	–58 58 54	0	UT
Orion BN	05 35 14.5	–05 22 22.0	I	UT + AT <sup>b</sup>
Orion SC3	05 35 16.2	–05 23 22.2	I/II	UT
IRAS15411–5352	15 44 59.503	–54 02 19.22	I	AT
IRAS17149–3916	17 18 23.9	–39 19 10	I	AT
IRAS17216–3801	17 25 06.3	–38 04 02	I/II	AT
IRAS18060–2005	18 08 58.3	–20 05 14	I	AT
IRAS18064–2020	18 09 21.0	–20 19 31	I	AT
M17SW IRS1	18 20 19.7	–16 13 31	I	AT

<sup>a</sup> See abstract for introduction.

<sup>b</sup> Snapshot with UTs, subsequent detailed studies at ATs.

Table 1: Target list sorted by decreasing science priority.



**MIDI guaranteed time observations –  
Abstract and source list**

**The dusty circumstellar environment of Hot Stars**

**Authors:** O. Chesneau (coordinator), L.B.F.M. Waters, B. Lopez, M. Robberto, F. Vakili,  
E. Bakker, A. Glazeborg-Kluttig, J. de Jong, S.Ligori

**1 Abstract**

The stellar winds of hot stars strongly affect the galactic interstellar medium, by many aspects of these winds are still poorly known. The winds of massive hot stars are driven by the strong radiation field, whose interaction with matter has been modeled in 1975 by Castor, Abbott and Klein in the frame of an isotropic and homogeneous wind. Since then, however, many observations have demonstrated that most of these winds are rather time variable, asymmetric, and/or clumpy. The combined effects of rotation, magnetic fields, pulsations and binarity certainly complicate the picture. Most remarkable is the fact that several classes of hot stars have large amounts of circumstellar dust.

It is likely that the presence of a dusty environment is closely related to the stellar wind. The conditions that lead to dust formation in that somewhat hostile environment are not well understood and require the knowledge of the dust spatial distribution. MIDI at the VLT Interferometer will be able to resolve the location and geometry of the dusty circumstellar environments of some bright hot stars ( $N < 1$ ), and to reveal the innermost regions where dust survives ( $\geq 100R_*$ ). This proposal addresses three topics related to hot stars with poorly understood circumstellar dust. The effects of binarity and stellar rotation on the presence of dust near hot stars will also be addressed.

The first two proposals deal with dust and gas in the very dense winds of evolved, very massive stars: (i) the enigmatic LBV  $\eta$  Car, and (ii) the hydrogen-poor Wolf-Rayet (WR) stars. In each case, binarity is suspected to be at the origin of the unexplained level of hot dust around these systems. The third project addresses the circumstellar environment of the B[e] stars. These are hot stars with large amounts of circumstellar gas and dust, whose origin, composition and geometry are poorly understood, but may be related to rapid rotation of the central star.

Titles of the subproposals:

- 'The geometry of the core of the  $\eta$  Carinae system'
- 'Hostile environnement around WC stars'
- 'B[e] stars dusty environnement'

Total observing time : **2 hours on UTs - used up**  
**68 hours on ATs**

## 2 Source list

⇒ these six objects are protected for observations on ATs only

Object name	Priority	$\alpha(2000)$	$\delta(2000)$	IRAS 12 $\mu\text{m}$ in Jy	Sp. Type	Telescopes
WR 98a	1	17 41 13.3	-30 32 33	51.6	WC8+?	ATs
HD 45677	1	06 28 17.42	-13 03 11.1	146	unclB[e]	ATs
HD 50138	1	06 51 33.40	-06 57 59.4	70.3	unclB[e]	ATs
HD 62623	1	07 43 48.47	-28 57 17.4	258	A2Iabe	ATs
MWC 300	2	18 29 25.68	-06 04 37.2	100.6	unclB[e]	ATs
CPD -42 11721	3	16 59 06.9	-42 42 08	95	unclB[e]	ATs

**Note:** Priorities range from 1 (high) to 3 (low)

## **MIDI guaranteed time observations –**

### **Abstract and source list**

## **Cool Late Type Stars and related objects**

Authors: A. Quirrenbach (coordinator), B. Lopez, E. Bakker, O. Chesneau, W.D. Cotton, Ch. Leinert, J. Meisner, J.-L. Menut, G. Niccolini, G. Perrin, P. Schuller, R. Tubbs, L.B.F.M. Waters

### **1 Abstract**

We propose to use MIDI at the VLTI to study mass loss in a sample of Asymptotic Giant Branch (AGB) stars, Red Supergiants (RSG), and related cool evolved objects with a dusty circumstellar environment. Both AGB stars and RSG are characterized by a dense, dusty stellar wind which is probably driven by stellar pulsations in combination with radiation pressure on dust. We wish to investigate the structure of these winds, in particular the location of the dust formation layer, and its dependence on pulsation cycle, mass loss rate and chemistry. Spectrally resolved interferometry in the 10  $\mu\text{m}$  window will allow a measurement of the stellar radius and/or the warm molecular layer (even in stars with high mass loss rates), and of the location of the different types of dust that condense in AGB outflows. Deviations from spherical symmetry in the inner regions of the wind and the effects of binarity on the geometry of the envelope will be investigated.

Total observing time : **25 hours on UTs - used up**  
**450 hours on ATs**

## 2 Source list

⇒ all these 28 objects are protected for observations on ATs only

target name	coordinates ( $\alpha, \delta$ ) (J2000)		telescopes	comments
IK Tau	03 53 28.6	+11 24 20	ATs	M6e
SW Vir	13 14 04.38	-02 48 25.1	ATs	M
R Hya	13 29 42.78	-23 16 52.8	ATs	Me
IRC+20326	17 31 55.0	+17 45 20	ATs	M2
U Her	16 25 47.47	+18 53 32.86	ATs	O-rich LPV
R Dor	04 36 45.59	-62 04 37.8	ATs	O-rich AGB low $\dot{M}$
Mira	02 19 20.79	-02 58 39.5	ATs	O-rich AGB moderate $\dot{M}$
R Leo	09 47 33.49	+11 25 43.6	ATs	O-rich AGB moderate $\dot{M}$
AFGL 5259	09 45 15.2	-22 01 45	ATs	O-rich AGB
R Aql	19 06 22.25	+08 13 48.0	ATs	Me
R Aqr	23 43 49.46	-15 17 04.20	ATs	Me+, Symbiotic LPV
W Hya	13 49 02.00	-28 22 03.5	ATs	O-rich AGB low $\dot{M}$
$\pi$ 1 Gru	22 22 44.21	-45 56 52.6	ATs	S, Tc
V Aql	19 04 24.16	-05 41 05.4	ATs	C
U Hya	10 37 33.27	-13 23 04.35	ATs	C-rich LPV
R Lep	04 59 36.35	-14 48 22.5	ATs	CHe
V Hya	10 51 37.25	-21 15 00.3	ATs	C
T Sgr	19 16 14.44	-16 58 17.06	ATs	C and Li rich LPV
R Scl	01 26 58.09	-32 32 35.5	ATs	C-rich AGB low $\dot{M}$
IRC +10 216	09 47 57.3	+13 16 44	ATs	C-rich AGB high $\dot{M}$
IRAS 15194-5115	15 23 05.0	-51 25 59	ATs	C-rich AGB high $\dot{M}$
IRAS 09116-2439	09 13 54.1	-24 51 21	ATs	(SiC)
VX Sgr	18 08 04.04	-22 13 26.61	ATs	RSG with disk
VY CMa	07 22 58.33	-25 46 03.2	ATs	RSG
$\alpha$ Ori	05 55 10.31	+07 24 25.4	ATs	RSG
Red Rectangle	06 19 58.22	-10 38 14.7	ATs	Binary post-AGB star
HR 4049	10 18 07.59	-28 59 31.2	ATs	Binary post-AGB star
RY Sgr	19 16 33	-33 31 20	ATs	R CrB star

**MIDI guaranteed time observations –  
Abstract and source list**

**Extra-solar planets and brown dwarfs**

Authors: Bruno Lopez (coordinator), Thomas Henning, Christoph Leinert,  
Jeffrey Meisner, Peter Schuller, Martin Vannier  
– in coordination with Romain Petrov, representing the AMBER programme

**1 Abstract**

Since the first detection of a hot giant planet orbiting around the star 51 Peg, several very exciting results have been obtained in detecting planets outside our solar system. We propose to observe, within the guaranteed time of VLTI/MIDI, two extrasolar planets (known hot Jupiter-like sources) and two brown dwarfs. The aim of the observations is to directly detect planet and brown dwarf radiation and to deduce spectral information of their atmospheres. Acquisitions of calibrated visibility curves and differential interferometry methods will be performed for reaching the best accuracy in the observations.

The observations of brown dwarfs will be compared to the observations of ATsthe Jupiter-like planets. One of our goals, for instance, will be to verify that the companion of GL86 is indeed a planet and not a brown dwarf itself.

Total observing time : **25 hours on UTs**

**2 Source list**

⇒ **these four objects are protected for UT observing only**

Object	$\alpha$	$\delta$	Sp. Type	V	N	Dist.	Msin(i)
GL 86	02 10 14.4	-50 50 00.5	F6	6.17	4	10.9 pc	4. $M_{Jup}$
GL 876	22 53 16	-14 15 44.3	M4	10.1	4.7	4.7 pc	0.5 $M_{Jup}$
HD 217580	23 01 51.54	-03 50 55.4	K4v	7.46	4.8	16.9 pc	60 $M_{Jup}$
HD 112758	12 59 01.56	-09 50 02.7	K0v	7.56	5.6	21.0 pc	35 $M_{Jup}$